



A project of Volunteers in Asia

Agro-Forestry Systems for the Humid Tropics East  
of the Andes

by: John P. Bishop

Published by:

Instituto Nacional de Investigaciones  
Agropecuarias  
Estacion Experimental Napo  
Centro Amazonico Limoncocha  
Apartado 2600  
Quito, Ecuador

Available for free if readers send payment to  
cover postage or wish to exchange publications.

Available from:

John P. Bishop  
Instituto Nacional de Investigaciones  
Agropecuarias  
Estacion Experimental Napo  
Centro Amazonico Limoncocha  
Apartado 2600  
Quito, Ecuador

Reproduced by permission of the U.S. Agency for  
International Development.

Reproduction of this microfiche document in any  
form is subject to the same restrictions as those  
of the original document.



AGRO-FORESTRY SYSTEMS  
FOR THE HUMID TROPICS

EAST OF THE ANDES

by

John P. Bishop

ESTACION EXPERIMENTAL "NAPO"

INSTITUTO NACIONAL DE INVESTIGACIONES AGROPECUARIAS

AGRO-FORESTRY SYSTEMS FOR THE HUMID TROPICS  
EAST OF THE ANDES: I. Integrated Foodcrop,  
Swine, Chicken and Fuelwood Production.\*

by

John P. Bishop\*\*

Estación Experimental Napo/Centro Amazónico Limoncocha\*\*\*  
Instituto Nacional de Investigaciones Agropecuarias  
Apartado 2600 - Quito, Ecuador

Introduction

Each region of the developing world has unique nutritional-demographic problems and potentials. Such problems are compounded in the central Andean region because its population has the highest density and birth rate in South America while consumption levels of protein and calories are among the lowest in the Western Hemisphere. Traditionally most people of Colombia, Ecuador, Perú and Bolivia have lived in highland areas.

These countries also contain vast humid tropical lowlands east of the Andes (Figure 1) into which highways are being built due to petroleum discoveries and colonization plans. Such highways are leading the poorest of the rural poor away from over-populated highland areas to establish small farms in the eastern lowlands (8). A sizeable part of each family farm is utilized for products of primary need such as foodcrops, swine, chickens and fuelwood.

---

\* Paper presented to International Conference on Amazonian Agricultural and Land Use Development, sponsored by ICRAF/CIAT/RF/GTZ/NCSU, Cali, Colombia, April 16-18, 1980. English translation of Spanish manuscript.

\*\* Member of INIAP/UFLA/IBRD Technical Assistance Mission. Author acknowledges assistance given by Summer Institute of Linguistics.

\*\*\* Average annual precipitation 3102 mm, elevation 243 masl, latitude 0°24'S.

A shifting form of agriculture is practiced by small farmers east of the Andes (25,26,32). As human populations and expectations increase, rest periods in shifting cultivation are shortened, accelerating in an alarming rate soil deterioration and weed/pest invasion, and critically decreasing yields precisely as needs are increasing (25, 30, 32).

One promising solution is to intensify the rest period with swine, chicken and fuelwood production (1,2,3,4,6,17, 19,21,28). Forage and fuelwood legumes increase soil aeration, organic matter, nitrogen and available phosphorus, as well as control soil erosion and leaching (4,5,20,27). Swine also improve soil fertility by depositing organic matter which stimulates legume/Rhizobium symbiosis and by supplying fecal microorganisms which mineralize crop residues (4). In addition, swine can improve small farm income (7) and produce low cost animal protein without cereal grains (29).

The American lowland tropics have the highest per capita swine population in the world (Table 1), more than five times that of tropical Africa and Asia (33). Most swine and chickens in the tropical lowlands east of the Andes are produced on small farms (Table 2) utilizing an open range production system with banana as the principle swine feed and corn as the supplemental chicken feed.

#### Integrated Foodcrop, Swine, Chicken and Fuelwood Production

In Amazonian Ecuador, studies are being realized to intensify open range swine husbandry utilizing the following mixture of perennial species in a multi-strata production system: Desmodium ovalifolium (trébol tropical), Canna edulis (achira), Musa acuminata x M. balbisiana ABB (orito) and Inga edulis (guaba).

The umbraphilus legume Desmodium ovalifolium constitutes the ground floor (19) as forage legumes are more palatable

and more efficiently utilized by swine than are grasses (10,14). The root forage Canna edulis and banana Musa acuminata x M. balbisiana ABB are local perennial crops with low labor and soil fertility requirements and produce low cost feeds for direct consumption by swine (11,15,16, 18,24,31,33). The fast growing native legume tree Inga edulis improves soil fertility and structure (22) as well as produces fuelwood (1,2,3) on an eight year rotation cycle (Figure 3).

Initially, conventional agriculture is practiced on a new plot each year: land clearing and production of short cycle species in conformity with one or another classical multiple cropping system. The perennial species of the future multi-strata system are interplanted during short cycle cropping (Figure 4). Following a two year period of short cycle cropping, the mentioned perennial components will have reached a vertically stratified state of development. Four distinct strata are rapidly differentiated and the resultant multi-strata structure and multi-species composition ecologically and biologically approaches a sustained yield forest ecosystem (9,12,13).

A ten hectare family unit (Figure 5) is divided into eight lots (1 ha. ea.) which are used following field crops (Table 3) for weanling pigs, fattening pigs and breeding sows. Also, eight lots (0.2 ha. ea.) are formed and used following garden crops (Table 4) for individual farrowing pens. Six strands of barbed wire and closely woven living Jatropha curcas (piñon) posts are used as fencing (23). Internal swine parasites are chemically controlled (levamisole) every three months synchronized with alternate grazing. Also, a 0.4 ha. lot is used for the farm home, chicken house and fruit trees (Table 5).




Figure 1. Dry months in the tropical lowlands east of the Andes  
(SOURCE: 1978 CIAT Annual Report).




Figure 2. Number of farms by size in Amazonian Ecuador  
(SOURCE: Ministerio de Agricultura y Ganadería, 1978).




Figure 3. Integrated foodcrop, swine and fuelwood production.





Figure 4. Average precipitation (15 years) and cropping practices in the Centro Amazónico Limoncocha




Figure 5. Integrated foodcrop, swine, chicken and fuelwood production in a 10 ha family unit.

---

Table 1. American countries with large swine populations  
(SOURCE: World Almanac, 1978).

---

	<u>Number of people per hog</u>
1. Brazil	2.3
2. Ecuador	2.7
3. Haiti	2.8
4. Nicaragua	3.3
5. U.S.A.	3.9

---

Table 2. Percentage of swine by farm size in Ecuador  
(SOURCE: Ministerio de Agricultura y Ganadería  
1978).

Farm size	% of swine
0 - 20 ha.	81.7 %
20 - 50 ha.	10.7 %
+ 50 ha.	7.6%

---

Table 3. Major foodcrops in Amazonian Ecuador.

---

<u>Local name</u>	<u>Scientific name</u>	<u>Variety</u>
Maíz	<u>Zea mays</u>	INIAP VS-2
Yuca	<u>Manihot esculenta</u>	Native
Plátano	<u>Musa acuminata x</u> <u>M. balbisiana AAB</u>	Local
Papa mandi	<u>Xanthosoma sagittifolium</u>	Native
Papa china	<u>Colocasia esculenta</u>	Local
Papaya	<u>Carica papaya</u>	Native

---

Table 4. Minor foodcrops in Amazonian Ecuador

<u>Local name</u>	<u>Scientific name</u>	<u>Variety</u>
Maní	<u>Arachis hypogaea</u>	Native
Fréjol común	<u>Phaseolus vulgaris</u>	Local
Fréjol ratón	<u>Vigna unguiculata</u>	Local
Fréjol vainita	<u>Vigna sesquipedalis</u>	Local
Habas nativas	<u>Phaseolus lunatus</u>	Native
Haba blanca	<u>Canavalia ensiformis</u>	Local
Ashipa	<u>Pachyrhizus tuberosus</u>	Native
Piña	<u>Ananas comosus</u>	Native
Cocona	<u>Solanum toapiro</u>	Native
Badea	<u>Passiflora quadrangularis</u>	Native
Granadilla	<u>Passiflora edulis</u>	Native
Maíz pequeño	<u>Zea mays</u>	Local
Camote	<u>Ipomea batatas</u>	Native
Papa de soga	<u>Dioscorea trifida</u>	Native
Pujín	<u>Calathea allouia</u>	Native
Achocha	<u>Cyclantera pedata</u>	Native
Tomate criollo	<u>Lycopersicon esculentum</u>	Local
Zapallo	<u>Cucurbita sp.</u>	Native
Cuchicol	<u>Alternanthera sp.</u>	Native
Cebolla criolla	<u>Allium capa</u>	Local
Caña de azúcar	<u>Sacharum sp.</u>	Local

Table 5. Fruit trees in Amazonian Ecuador

<u>Local name</u>	<u>Scientific name</u>	<u>Variety</u>
Limón mandarina	<u>Citrus limonia</u>	Local
Lima	<u>Citrus limettoides</u>	Local
Naranja criollo	<u>Citrus sinensis</u>	Local
Maní de árbol	<u>Caryodendron orinocensis</u>	Native
Guaba Ilta	<u>Inga densiflora</u>	Native
Arbol de pan	<u>Artocarpus alfilis</u>	Local
Cacao blanco	<u>Theobroma bicolor</u>	Native
Zapote	<u>Calocarpum sapote</u>	Native
Abiyu	<u>Pouteria cecropiaeefolia</u>	Native
Anona	<u>Annona squamosa</u>	Native
Uvilla	<u>Pourouma cecropiaeefolia</u>	Native
Guaba común	<u>Inga edulis</u>	Native
Guaba machetona	<u>Inga spectabilis</u>	Native
Aguacate	<u>Persea americana</u>	Native
Guanábana	<u>Annona muricata</u>	Native
Chonta duro	<u>Guilielma gasipaes</u>	Native
Guayaba	<u>Psidium guajava</u>	Native

Literature Cited

1. Bishop, J. P. 1978. The development of a sustained yield agro-ecosystem in the upper Amazon. *Agro-Ecosystems*, 4: 459-461.
2. Bishop, J. P. 1978. Desarrollo y transferencia de tecnología para pequeñas fincas en la Región Amazónica ecuatoriana. En: Seminario sobre Manejo de los Sistemas Ecológicos y Alternativas de Producción Agro-Silvo-Pastoril en la Región Amazónica Ecuatoriana, patrocinada por el Instituto Nacional de Colonización de la Región Amazónica Ecuatoriana, Limoncocha, Ecuador. 8 pp.
3. Bishop, J. P. 1979. Producción familiar agro-porcino-forestal en el trópico húmedo hispanoamericano. En: Taller sobre sistemas Agro-Forestales en América Latina Tropical, patrocinada por CATIE/UNU, Turrialba, Costa Rica. 9 pp.
4. Bredero, T. J. 1977. The role of farmyard manures and green manures in soil fertility restoration in the humid tropics. *Abstr. Trop. Agric.* 3: 9-17.
5. Bredero, T. J. 1973. Green manuring and the N and P supply of swamp rice. *Nigerian Agric. J.*, 10: 248-257.
6. Breitenbach, C. A. 1974. Farming systems for the tropics and subtropics. In: *Guide for Field Crops in the Tropics and the Subtropics*. USAID, Washington, D. C., 22-28 pp.
7. CIAT, 1971. Sistemas de producción de ganado porcino. En: Informe Anual. Centro Internacional de Agricultura Tropical, Cali, Colombia. 41-52 pp.
8. Crist, R. E. and Nissly, C. M. 1973. East from the Andes. Univ. Florida Social Sciences Monograph No. 50. Univ. Florida Press, Gainesville, Florida. 166 pp.
9. Dubois, J. 1977. Investigaciones sobre Trópico Húmedo Americano. En: Seminario sobre Ecología del Trópico Húmedo Americano, IICA-TROPICOS, Mérida, Venezuela. IX AI-IX AIO pp.
10. Eyles, D. E. 1963. Integration of pigs into grassland farming. In: *Animal Health, Production and Pasture*. Longmans, London, 359-383 pp.

11. Herklots, G. A. 1972. Vegetables in South-East Asia. Hafner, New York. 525 pp.
12. Holdridge, L. R. 1959. Ecological indications of the need for a new approach to tropical land-use. Economic Botany, 13: 271-280.
13. Janzen, D. H. 1973. Tropical agroecosystems. Science, 182: 1212-1219.
14. Jones, D. W. and Wallace, H. D. 1974. Grain and forage crops for swine. In: Swine Production in Florida. Florida Department of Agriculture, Bull. No. 21, 93-99 pp.
15. Kay, D. E. 1973. Queensland Arrowroot. In: TPI Crop and Product Digest, No. 2 Root Crops, Tropical Products Institute, Overseas Development Administration, London. 120-126 pp.
16. Kurita, K. 1967. The cultivation of Canna edulis, and its value as a feedcrop. Japan J. Trop. Agric., 11 (1-2): 5-8.
17. Kirby, J. M. 1976. Agricultural land-use and the settlement of Amazonia. Pacific Viewpoint, 15: 105-131.
18. LeDividich, J. 1977. Feeding value of Canna edulis roots for pigs. J. Agric. Univ. P. R. 61 (3): 267-274.
19. Masefield, G. B. 1965. A Handbook of Tropical Agriculture. Oxford, London, 196 pp.
20. Moore, A. W. 1967. Changes in soil moisture and organic matter under different covers at Ibadan, Nigeria. Plant Soil, 27: 463-467.
21. Nye, P. H. and Greenland, D. J. 1960. The Soil under Shifting Cultivation. Commonwealth Bur. Soils, Tech. Commun. 51, Harpenden, U. K. 156 pp.
22. Ochse, J. J., et. al. 1961. Tropical and Subtropical Agriculture, Vol. 1. Macmillan, New York, 760 pp.
23. Payne, W. J. 1973. Disposición y Manejo de fincas tropicales. En: Ganadería en los Trópicos. Asociación Venezolana de Criadores de Ganado Cebú. Caracas, Venezuela. Vol. 1, 563 pp.
24. Purseglove, J. W. 1972. Tropical Crops: Monocotyledons. Wiley, New York. 607 pp.

25. Sánchez, P. A. 1977. Alternativas al sistema de agricultura migratoria en América Latina. En: Reunión sobre Manejo, Conservación de Suelos y Agricultura Migratoria en América Latina, FAO/SIDA, Lima, Perú. 44 pp.
26. Sánchez, P. A. 1973. Manejo de Suelos bajo sistemas de roza. En: Un Resumen de las Investigaciones Edafológicas en la América Latina Tropical. N. C. Agr. Exp. Sta. Tech. Bull. 219: 51-74.
27. Singh, A. 1967. Long-term effects of green manures in sub-tropics. Indian J. Agric. Sci., 37: 226-233.
28. Sprague, H. B. 1976. Combined crop/livestock farming systems for developing countries of the tropics and sub-tropics. Technical Series Bulletin No. 19, Office of Agriculture, Technical Assistance Bureau, Agency for International Development, Washington D. C., 30 pp.
29. Thomsen, M. 1978. The Farm on the River of Esmeralds. Mifflin, Boston. 329 pp.
30. Tosi, J. 1974. Desarrollo forestal del trópico americano frente a otras actividades económicas. En: Reunión Internacional sobre Sistemas de Producción para el Trópico Americano. Informes sobre Cursos, Conferencias y Reuniones No. 41, IICA. Zona Andina, Lima, Perú. 13 pp.
31. Walker, R. H. 1953. Some notes on the edible canna and its use in feeding pigs on the Lehmann system. Govt. Kenya; Report of an enquiry into the general economy of farming in the highlands. pp. 56-57.
32. Watters, W. F. 1971. Shifting cultivation in Latin America. FAO Forestry Dev. Paper No. 17. 305 pp.
33. Williamson, G. and Pyne, W. J. 1975. La Ganadería en Regiones Tropicales. Editorial Blume, Barcelona. 468 pp.

AGRO-FORESTRY SYSTEMS FOR THE HUMID TROPICS

EAST OF THE ANDES:

II. Integrated Cattle and Timber Production.\*

by

John P. Bishop\*\*

Estación Experimental Napo/Centro Amazónico Limoncocha\*\*\*

Instituto Nacional de Investigaciones Agropecuarias

Apartado 2600 - Quito, Ecuador

Introduction

The humid tropics east of the Andes are presently undergoing large scale deforestation in favor of cattle development. Forests are substituted with pastures following short-term cropping. Such development has provoked severe criticism as there occurs serious soil deterioration and few pastures persist. Today many of these pastures are found abandoned (1, 9, 13).

One suggested solution is to associate forage grasses with forage legumes and timber trees (2,3,4,5,6,7,8,10,11, 12,14,15,16). Forage legumes and timber trees can fulfill the following functions: a) significantly increase soil nitrogen by root associations with bacteria and fungi, b) significantly fertilize soil through leaf-fall, c) notably improve soil texture and aeration by physical and chemical effects, and d) substantially increase income from small farm pastures by sale of timber.

---

\* Paper presented to International Conference on Amazonian Agricultural and Land Use Development, sponsored by ICRAF/CIAT/RF/GTZ/NCSU, Cali, Colombia, April 16-18, 1980. English translation of Spanish manuscript.

\*\* Member of INIAP/UFLA/IBRD Technical Assistance Mission. Author acknowledges assistance given by Summer Institute of Linguistics.

\*\*\* Average annual precipitation 3102 mm, elevation 243 masl, latitude 0°24'S.

### Integrated Cattle and Timber Production

In Amazonian Ecuador, studies are being realized to evaluate the forage grass Brachiaria humidicola (kikuyo amazónico), legume Desmodium ovalifolium (trébol tropical) and timber tree Cordia alliodora (laurel) in a silvo-pastoral system (Figure 1).

At the beginning of the rainy season the B. humidicola and D. ovalifolium are established using vegetative material and planting stick. The C. alliodora is also transplanted (400/ha) at this time using rootstumps (Figures 2 & 3).

The newly established pastures are not grazed for one year or until timber trees are three meters high. Two years after reforestation, trees are thinned to 200/ha and again after four years leaving 100 high-grade trees per hectare (Figure 4).

### The Benefit

One hectare of pasture maintaining two bovines with 25% extraction per year will produce ten bovines in twenty years. Estimating each adult bovine at US \$300, cattle income per hectare in twenty years will be US \$3000.

One hundred Cordia trees per hectare can produce 100 m<sup>3</sup> of timber in twenty years. Estimating each m<sup>3</sup> of Cordia at US \$30, forest income per hectare in twenty years would be US \$3000.

Therefore, timber production has great potential to improve the economic productivity, ecological stability and sociological viability of small cattle farms in the humid tropics east of the Andes.

For technology transfer to the rural masses, small farmer training materials are being prepared for use in local adult education classes, regional radio education courses and practical classes in rural schools.




Figure 1. Integrated cattle and timber production.




Figure 2. Average precipitation (15 years) and planting sequence in the Centro Amazónico Limoncocha.




Figure 3. Planting diagram.




Figure 4. Mixed production system for 50 ha farms.

### Literature Cited

1. Anon. 1978. Forestry for local community development. FAO Forestry Paper No. 7, Rome, 114 pp.
2. Bishop, J. P. 1978. The development of a sustained yield tropical agro-ecosystem in the upper Amazon. *Agro-Ecosystems*, 4: 459-461.
3. Bishop, J. P. 1978. Desarrollo y transferencia de tecnología para pequeñas fincas en la Región Amazónica Ecuatoriana. En: Seminario sobre Manejo de los Sistemas Ecológicos y Alternativas de Producción Agro-Silvo-Pastoril en la Región Amazónica Ecuatoriana, patrocinada por el Instituto Nacional de Colonización de la Región Amazónica Ecuatoriana, Limoncocha, Ecuador.
4. Bishop, J. P. 1979. Producción ganadera-forestal en el trópico húmedo hispanoamericano. En: XIII Conferencia Anual sobre Ganadería y Avicultura en América Latina, patrocinada por la Universidad de Florida, Gainesville.
5. Bishop J. P. and Muñoz, K. 1979. Producción ganadera-forestal en la Región Amazónica Ecuatoriana. En: Reunión de Trabajo sobre Pastos Tropicales, patrocinada por el Centro Internacional de Agricultura Tropical, Cali, Colombia.
6. Cook, B. G. and Grimes, R. F. 1977. Multiple land-use of open forest in south-eastern Queensland for timber and improved pasture: Establishment and early growth. *Trop. Grasslands*, 11:239-245.
7. Gregory, E. W. 1972. Integration of grazing in tropical forestry---An experiment in combining cattle raising with pine plantation forestry in Fiji. Seventh World Forest Congress, Buenos Aires.
8. Kennard, D. E. and Walker, B. H. 1973. Relationships between tree canopy cover and *Panicum maximum* in the vicinity of Fort Victoria. Rhodesian J. Agric. Res., 11: 145-153.
9. King, K. F. S. and Chandler, M. T. 1978. The wasted lands. International Council for Research in Agro-forestry, Nairobi.
10. Kirby, J. M. 1976. Agricultural land-use and the settlement of Amazonia. *Pacific Viewpoint*, 15: 105-131.
11. Kirby, J.M. 1976. Forest grazing: A technique for the tropics. *World Crops*, 28: 248-251.

12. Knowles, R. L., Klomp, B. K. and Gillingham, A. 1977. Report for the Fiji Pine Commission on forest grazing research. Rotorua. New Zealand Forest Service, 13 p.
13. Parson, J.J. 1976. Forest to pasture: Development or destruction? Rev. Bio. Trop., 24 (Supl. 1): 121-138.
14. Payne, W. J. 1976. Possibilities for the integration of tree crops and livestock production in the wet tropics. J. Sci. Food Agric., 27:888.
15. Peck, R. B. 1977. Sistemas agro-silvo-pastoriles como una alternativa para la reforestación en los trópicos americanos. Bogotá, Colombia, CONIF, 73-84 p.
16. Thomas, D. 1978. Pastures and livestock under tree crops in the humid tropics. Trop. Agric. (Trinidad) 55: 39-44.
17. Wyatt-Smith, J. 1979. Agro-forestry in the Tropics: A new emphasis in rural development. Span 22: 65-67.